

## How Will Labor Be Different If I Choose Labor Induction at 39 Weeks?

The American College of Obstetricians and Gynecologists now states that it is reasonable to offer induction at 39 weeks to women with low-risk pregnancies who are sure of their due dates.<sup>1</sup> Induction at 39 weeks does not result in better outcomes for the baby, but it may *slightly* reduce the risk for cesarean surgery. If you choose induction at 39 weeks, your experience of labor and birth will be different than if you let labor begin on its own. You will require more routine medical interventions. This chart explains the differences. It is based on the most common labor induction method of using IV synthetic oxytocin (known as Pitocin in the U.S. and in Canada) to induce labor and assumes that your pregnancy is considered low-risk.

INDUCED LABOR AT 39 WEEKS	LABOR BEGINS ON ITS OWN
<b>CERVICAL RIPENING</b>	
<p>Usually some time during the 38th week of pregnancy, your health care provider will do a vaginal exam and evaluate the readiness of your cervix for birth using the “Bishop Score.” If your modified Bishop Score is less than 5, indicating that your cervix is not ready for labor, you will undergo “cervical ripening” in conjunction with or followed by IV administration of synthetic oxytocin.<sup>1,2</sup> Cervical ripening will be one of the first procedures done after you have been admitted to the hospital for induction or sometimes it is done the day or evening before your scheduled induction. Medications or devices may be used to “ripen” your cervix so that it will stretch (dilate) for labor.</p>	<p>In the weeks and days before labor starts on its own, your cervix will gradually soften, thin, and may begin to open (dilate). In addition to cervical ripening, hormones (catecholamines) are released in the weeks or days before labor begins to prepare the fetal lungs for air breathing.<sup>3,4</sup></p>
<b>IV FLUIDS</b>	
<p>Yes. You will have IV fluids, including IV synthetic oxytocin (Pitocin).</p>	<p>No. You will be encouraged to drink fluids to meet hydration and caloric needs.<sup>5</sup></p>
<b>FETAL MONITORING</b>	
<p>You will have continuous electronic fetal monitoring.<sup>6</sup> The monitors can be external (2 belts) or internal.</p>	<p>The baby’s heart beat can be monitored intermittently with a hand-held device<sup>5</sup> – usually every 15 to 30 minutes during active labor and every 15 minutes in 2nd stage before pushing and every 5 minutes during pushing.<sup>7</sup></p>
<b>MOBILITY/AMBULATION DURING LABOR</b>	
<p>Most laboring women who are induced with IV synthetic oxytocin stay in bed or stand or sit right next to the bed. A few women walk with portable monitoring devices called telemetry.</p>	<p>You are free to move around, choosing the positions most comfortable for you, and walking if you wish.<sup>5</sup></p>
<b>LABOR CONTRACTION PAIN/PAIN MANAGEMENT</b>	
<p>Your contractions may peak sooner and feel more intense than spontaneous labor contractions. You are more likely to ask for epidural analgesia.<sup>8</sup></p>	<p>You can try nonpharmacologic comfort measures such as a warm bath or shower.<sup>5</sup> (Note: Epidural analgesia requires IV fluids and continuous fetal monitoring.)</p>
<b>NEED FOR URINARY CATHETER</b>	
<p>If you have epidural analgesia, you may have problems urinating.<sup>9</sup> You may need to be catheterized to empty your bladder.</p>	<p>Problems with urinating are associated with epidural analgesia.</p>
<b>LENGTH OF LABOR</b>	
<p>Your labor may be longer than if labor had begun on its own. Health care providers are encouraged to allow up to 24 hours or more for the latent phase of labor and to administer Pitocin for at least 12-18 hours after membrane rupture before calling the induction a failure (and recommending cesarean surgery).<sup>10</sup></p>	<p>The latent phase of labor can be prolonged (&gt; than 20 hours in first-time mothers). However, you are encouraged to stay at home for most of this time, only going to the hospital when labor becomes “active” at about 6 cm.<sup>5,10</sup></p>

## REFERENCES

1. ACOG & SMFM. (August 8, 2018). *Leaders in obstetric care respond to the published results of the ARRIVE Trial*. [www.acog.org/About-ACOG/News-Room/Statements/2018/Leaders-in-Obstetric-Care-Respond-to-the-Published-Results-of-the-ARRIVE-Trial](http://www.acog.org/About-ACOG/News-Room/Statements/2018/Leaders-in-Obstetric-Care-Respond-to-the-Published-Results-of-the-ARRIVE-Trial)
2. Grobman, W. A. (2018). Labor induction versus expectant management in low-risk nulliparous women (Supplementary appendix) [protocol for labor induction used in the ARRIVE Trial]. *New England Journal of Medicine*, 379(6), 513-523. <http://dx.doi.org/10.1056/NEJMoa1800566>
3. Jain, L. (2006). Implications of labor on neonatal outcomes. In *NIH State-of-the-Science Conference: Cesarean delivery on maternal request*. Retrieved 8/02/18 from <http://consensus.nih.gov/2006/cesareanabstracts.pdf>.
4. Jain, L. & Eaton, D.C. (2006). Physiology of fetal lung fluid clearance and the effect of labor. *Seminars in Perinatology*, 30(1), 34-43. <http://dx.doi.org/10.1053/j.semperi.2006.01.006>
5. ACOG. (2019). Committee opinion #766: Approaches to limit intervention during labor and birth. *Obstetrics & Gynecology*, 133(2), e164-e173. <https://bit.ly/2MXfjLw>
6. ACOG. (2009, reaffirmed 2019). ACOG practice bulletin #107: Induction of labor. *Obstetrics & Gynecology*, 114(2) Part 1, 386-397. <http://dx.doi.org/10.1097/AOG.0b013e3181b48ef5>
7. American College of Nurse-Midwives (ACNM). (2015). ACNM Clinical bulletin: Intermittent auscultation for intrapartum fetal heart rate surveillance. *Journal of Midwifery & Women's Health*, 60(5), 626-632. <http://dx.doi.org/10.1111/jmwh.12372>
8. Poma, S., et al. (2017). Outcomes of induced versus spontaneous labor. *Journal of Maternal-Fetal & Neonatal Medicine*, 30(10), 1133-1138. <http://dx.doi.org/10.1080/14767058.2016.1205028>
9. ACOG. (2019). Practice bulletin: Obstetric Analgesia and anesthesia. *Obstetrics & Gynecology*, 133(3), e208--e225. <http://dx.doi.org/10.1097/AOG.0000000000003132>
10. ACOG & SMFM. (2014). Obstetric care consensus: Safe prevention of the primary cesarean delivery. *Obstetrics & Gynecology*, 123(3), 693-711. <http://dx.doi.org/10.1097/01.AOG.0000444441.04111.1d>

### Also Recommended

- Amis, D. (2019). Healthy birth practice #1: Let labor begin on its own. *Journal of Perinatal Education*, 28(2), 68-80.



June 2019. Chart compiled by Debby Amis for The Family Way Publications, [www.thefamilyway.com](http://www.thefamilyway.com).

May be reproduced in full with attribution for noncommercial purposes.

Like us on facebook at [www.facebook.com/familywaypublications](http://www.facebook.com/familywaypublications) for new handouts and childbirth education research updates.