

The Bottom Line: When is Induction Appropriate?

References

1. American College of Obstetricians and Gynecologists (ACOG) and the Society for Maternal-Fetal Medicine (SMFM). (2018). *Leaders in obstetric care respond to the published results of the ARRIVE trial*. www.acog.org/patient-resources/faqs/labor-delivery-and-postpartum-care/induction-of-labor-at-39-weeks
2. World Health Organization (WHO). (2018). *WHO recommendations: Induction of labour at or beyond term* (page 5). Geneva, Switzerland. <https://apps.who.int/iris/bitstream/handle/10665/277233/9789241550413-eng.pdf?ua=1>
3. Society of Obstetricians and Gynaecologists of Canada. (2017). Practice guideline #214: Guidelines for the management of pregnancy at 41+0 to 42+0 weeks. *Journal of Obstetrics and Gynaecology Canada*, 39(8), e164-e174. www.ncbi.nlm.nih.gov/pubmed/28729108
4. Keulan, J. K.J., et al. (2019). Induction of labour at 41 weeks versus expectant management until 42 weeks (INDEX): multicentre, randomised non-inferiority trial. *BMJ*, 2019 Feb 20;364:l344. doi: 10.1136/bmj.l344. www.bmj.com/content/364/bmj.l344.long
5. Wennerholm, U-B., et al. (2019). Induction of labour at 41 weeks versus expectant management and induction of labour at 42 weeks (SWEdish Post-term Induction Study, SWEPIS): multicentre, open label, randomised, superiority trial. *BMJ*, 2019;367:l6131. www.bmj.com/content/367/bmj.l6131.long
6. Rydahl, E., Declercq, E., Juhl, M., & Maimburg, R.D. (2019). Routine induction in late-term pregnancies: follow-up of a Danish induction of labour paradigm. *BMJ Open*, 2019;9:e032815. doi:10.1136/bmjopen-2019-032815. <https://bmjopen.bmj.com/content/bmjopen/9/12/e032815.full.pdf>
7. Bergsjö, P., Denman III, D.W., Hoffman, H.J. & Meirik, O. (1990). Duration of human singleton pregnancy: A population-based study. *Acta Obstet Gynecol Scand*, 69(3), 197-207. www.ncbi.nlm.nih.gov/pubmed/2220340
8. Middleton, P., Shepherd, E. & Crowther, C.A. (2018). Induction of labour for improving birth outcomes for women at or beyond term. *Cochrane Database of Systematic Reviews 2018, issue 5*.art.no. CD004945. www.cochrane.org/CD004945/PREG_induction-labour-women-normal-pregnancies-or-beyond-term
9. Grobman, W. A., et al. (2018). Labor induction versus expectant management in low-risk nulliparous women. *New England Journal of Medicine*, 379(6), 513-523. www.nejm.org/doi/10.1056/NEJMoa1800566?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%3dwww.ncbi.nlm.nih.gov
10. Rydahl, E., Eriksen, L. & Juhl, M. (2018). Effects of induction of labor prior to post-term in low-risk pregnancies: a systematic review. *JBIC Database of Systematic Reviews and Implementation Reports*, 17(2):170-208. www.ncbi.nlm.nih.gov/pmc/articles/PMC6382053/
11. Derrick, D.C. & Higson, N. (2019). Labor induction at term – How great is the risk of refusing it? *AIMS Journal*, 31(9). 5-12. www.aims.org.uk/journal/item/induction-at-term

Additional Recommended Reading

Dekker, R. (2020). Evidence on: Inducing for due dates.

<https://evidencebasedbirth.com/evidence-on-inducing-labor-for-going-past-your-due-date/>



Updated May 2020. Chart compiled by Debby Amis for The Family Way Publications, www.thefamilyway.com.

May be reproduced in full with attribution for non-commercial purposes.

Like us on Facebook at www.facebook.com/familywaypublications for new handouts and childbirth education research updates.